



You received an immunotherapy medication called **Diphenylcyclopropenone (DPCP)** that was applied to your wart today (Only one wart if you have multiple). You can expect the site to become red, itchy and rashy. This is expected and a desired outcome.

Do not get wet or cover for 4 hours after the application

Care when you go home:

1. After 1 week you can begin using the product you have at home DPCP 1%. If tolerated you can apply to more than one of your warts if you have multiple. Wash hands.

- Use the end of a Qtip with the cotton tip broke off and dip the stick into the medication and apply to wart.
- Be careful not to get on normal skin.
- This will dry very quickly
- Wash your hands after applying your medication.
- Do not get wet or cover for 4 hours after the application

2. Apply the DPCP using above instructions once per week for 2 weeks on non consecutive nights

- Wash hands.
- Use the end of a Qtip with the cotton tip broke off and dip the stick into the medication and apply to wart.
- Be careful not to get on normal skin.
- This will dry very quickly
- Wash your hands after applying your medication.
- Do not get wet or cover for 4 hours after the application

3. After the 2 weeks, increase use of DPCP to 3 times per week on non-consecutive nights

- Wash hands.
- Use the end of a Qtip with the cotton tip broke off and dip the stick into the medication and apply to wart.
- Be careful not to get on normal skin.
- This will dry very quickly
- Wash your hands after applying your medication.
- Do not get wet or cover for 4 hours after the application.

Continue using until your next office visit with the physician.

Follow up with the MD in 3 mos